

PROGRESS IN NIOBIUM AND TANTALUM COORDINATION CHEMISTRY

Rolf W. Berg

Chemistry Department A, The Technical University of Denmark, DK-2800 Lyngby, Denmark

(Received September, 18 1990)

CONTENTS

| | | |
|-------|---|----|
| 1. | Introduction | 3 |
| 2. | Nb(V) and Ta(V) compounds (d^0) | 5 |
| 2.1 | Nb(V) and Ta(V) complexes containing only halides | 5 |
| 2.2 | Nb(V) and Ta(V) oxyhalide complexes | 10 |
| 2.2.1 | Oxyfluoride complexes | 10 |
| 2.2.2 | Other Oxyhalide complexes | 13 |
| 2.3 | Nb(V) and Ta(V) peroxyhalide and peroxy-pseudohalide complexes | 15 |
| 2.4 | Nb(V) and Ta(V) chalcogeno-halide complexes | 16 |
| 2.5 | Nb(V) and Ta(V) halide complexes with O-donor ligands | 17 |
| 2.5.1 | Fluoride complexes with O-donor ligands | 17 |
| 2.5.2 | Chloride complexes with O-donor ligands | 18 |
| 2.5.3 | Bromide complexes with O-donor ligands | 23 |
| 2.6 | Nb(V) and Ta(V) halide complexes with N-donor ligands | 23 |
| 2.7 | Nb(V) and Ta(V) oxyhalide complexes with N-donor ligands | 28 |
| 2.8 | Nb(V) and Ta(V) thiohalide complexes with N-donor ligands | 31 |
| 2.9 | Nb(V) and Ta(V) halide complexes with C-donor and other ligands | 32 |
| 2.10 | Nb(V) and Ta(V) pseudo-halide complexes with oxy-donors | 33 |
| 2.11 | Oxides of niobium(V) and tantalum(V) | 36 |
| 2.12 | Heteropolycompounds of Nb(V) and Ta(V) | 49 |
| 2.13 | Perchlorates, sulphates, phosphates and other Oxy-complexes of Nb(V) and Ta(V) | 50 |
| 2.14 | Intercalation of molecules into Nb(V) and Ta(V) oxides | 54 |
| 2.15 | Peroxy-niobates(V) and -tantalates(V) | 55 |
| 2.16 | Nb(V) and Ta(V) sulphides and selenides | 55 |
| 2.17 | Nb(V) and Ta(V) complexes with O-donor ligands | 57 |

| | | |
|--------|---|-----|
| 2.17.1 | Nb(V) and Ta(V) alkoxides | 57 |
| 2.17.2 | Nb(V) and Ta(V) alkenoxides | 62 |
| 2.17.3 | Nb(V) and Ta(V) carboxylates | 63 |
| 2.18 | Nb(V) and Ta(V) complexes with S- and Se-donor ligands | 64 |
| 2.19 | Nb(V) and Ta(V) complexes with O- and S-donor ligands | 66 |
| 2.20 | Nb(V) and Ta(V) oxynitrides and complexes with O- and N-donors | 68 |
| 2.21 | Nb(V) and Ta(V) complexes with As-ligands | 71 |
| 2.22 | Nb(V) and Ta(V) carbides and complexes with C-donor ligands | 71 |
| 2.23 | Biological complexes of Nb(V) and Ta(V) | 72 |
| 2.24 | Solvent extraction of Nb(V) and Ta(V) | 73 |
| 3. | Nb(V) and Nb(IV) mixed complexes (d^0 and d^1) | 74 |
| 4. | Nb(IV) and Ta(IV) compounds (d^1) | 75 |
| 4.1 | Nb(IV) and Ta(IV) halide complexes | 75 |
| 4.2 | Nb(IV) and Ta(IV) oxyhalide complexes | 77 |
| 4.3 | Nb(IV) and Ta(IV) halide complexes with O-donors | 77 |
| 4.4 | Other Nb(IV) and Ta(IV) halide complexes | 79 |
| 4.5 | Nb(IV) and Ta(IV) oxides | 80 |
| 4.6 | Nb(IV) and Ta(IV) complexes with O-donors | 80 |
| 4.7 | Nb(IV) and Ta(IV) sulphides and selenides | 81 |
| 4.8 | Other Nb(IV) and Ta(IV) complexes | 84 |
| 4.9 | Nb(IV) and Ta(IV) complexes with N- and C-donors | 84 |
| 5. | Mixed Nb(IV) and Nb(III) complexes (d^1 and d^2) | 85 |
| 6. | Nb(III) and Ta(III) complexes (d^2) | 85 |
| 6.1 | Nb(III) and Ta(III) halides and halide complexes | 85 |
| 6.2 | Nb(III) and Ta(III) chalcogenohalide complexes | 86 |
| 6.3 | Nb(III) and Ta(III) halides with O- and chalcogen-donors | 86 |
| 6.4 | Nb(III) and Ta(III) halides with N- and P-donors | 89 |
| 6.5 | Nb(III) and Ta(III) halides with C-donors | 91 |
| 6.6 | Nb(III) and Ta(III) oxides, sulphides and selenides | 94 |
| 6.7 | Nb(III) and Ta(III) nitrides | 96 |
| 6.8 | Nb(III) and Ta(III) C-donor complexes | 97 |
| 7. | Niobium and tantalum clusters of oxidation state < III | 97 |
| 8. | Nb(II) and Ta(II) complexes (d^3) | 102 |
| 9. | Nb(I) and Ta(I) complexes (d^4) | 102 |
| 10. | Nb(0) and Ta(0) complexes and alloys (d^5) | 104 |
| 11. | Niobium and tantalum in oxidation states below 0 | 105 |
| 12. | Compounds of unknown oxidation state | 106 |
| | References | 109 |

